

SYNTHESIS OF AMINOPICALINES WITH GOSSYPOL SCHIFF BASE AND PREPARATION OF METALLOCOMPLEXES WITH Cu^{2+} , Ni^{2+} SALTS

***Azizova Safina Isroil qizi
Khakberdiyev Shukhrat Makhramovich
Jizzakh Polytechnic Institute
E-mail: h.shyxrat81@gmail.com***

Abstract: Physicochemical properties of gossypol, their Schiff bases and metallocomplexes were synthesized. When synthesizing derivatives of Schiff's bases, gossypol and primary chiamin are taken in a 1:2 mol ratio, enough 96% ethyl alcohol is poured for its dissolution and stirred for three hours while heated in a magnetic stirrer. Structures of all obtained substances were analyzed using IR and UV spectra.

Key words: Gossypol, amino compound, Schiff's base, spectrum, biologically active substance, metallocomplex, polyphenol, triterpene, aldehyde, cotton plant, naphthalene.

Introduction

Gossypol is a specific polyphenolic compound and is considered the main polyphenolic compound of the cotton plant. belongs to the group of triterpene aldehydes and is found in plants called Gossipium. Among such plants, the cotton plant is included, in various parts of it, including leaves, roots, and other parts of the body in different amounts (up to 0,56-3%, depending on the type of cotton). occurs.

Gossypol is an amorphous, crystalline, lemon-yellow natural substance that is soluble in various organic solvents. In addition to the presence of various polar groups in the gossypol molecule, it is practically insoluble in water due to the presence of two isopropyl groups in the naphthalene residue.

Gossypol is very soluble in methyl, ethyl, isopropyl and butyl alcohols, as well as diethylene glycol, dioxane, acetone, diethyl ether, ethyl acetate, chloroform, carbon tetrachloride, dichloroethane, phenol, pyridine and heated naphthalene, heated cottonseed oil. In addition, gossypol is soluble in limited amounts in glycerol, cyclohexane, benzene, gasoline, and petroleum ether.

As a result of a number of studies, it was determined that the aldehyde groups in the gossypol ($C_{30}H_{30}O_8$) molecule form internal hydrogen bonds with the two oxygen atoms of the hydroxyl groups in the second -ortho position.



Substances formed as a result of the reaction of substances containing an amino group with gossypol are called Schiff's bases or azomethine derivatives. When synthesizing derivatives of Schiff's bases, gossypol and primary chiamin are taken in a 1:2 mol ratio, enough 96% ethyl alcohol is poured in to dissolve it, and it is stirred in a heated state (70-80 °C) in a magnetic stirrer for three hours, the progress of the reaction is monitored using YuQX. After the reaction is complete, the resulting substance is left for one day to settle completely, then it is filtered and washed 2-3 times with ethyl alcohol. The obtained substance is dried in a place where sunlight does not fall.

Table-1

Physicochemical quantities of synthesized Schiff bases

Synthesized Schiff base	Liquid °S	R _f	Color	Reaction product in (%)
Di-(2-amino-4-pikolin)gossipol	21 3-214	0 ,49	yellow	89
Di-(6-amino-3-pikolin)gossipol	20 9-210	0 ,44	yellow	92

Gossypol contains six –OH groups, two carbonyl, two methyl, two isopropyl and two naphthalene rings. In its IR-spectrum (n, cm-1) we can see the valence vibration frequencies belonging to the –OH group in the 3495, 3424 areas. The area between 1614 and 1441 indicates the presence of two naphthalene rings.

In the IR spectrum of the Schiff base formed by gossypol with 2-amino-4-picoline, we can see a clear change of the absorption maxima at 3240-3430 cm⁻¹ belonging to the NH₂ group.

When analyzing the IR spectrum, it can be observed that the valence vibrations at 1712 cm⁻¹ corresponding to the -CHO group have completely disappeared.

In the UV spectrum of the Schiff base of gossypol formed with 2-amino-4-picoline, a bathochromic shift occurs for this substance due to absorption in the 240,54 nm branch, and absorptions at 280, 427,98 nm are due to the hypsochromic shift of the Schiff base with N in the azomethine bond. is explained by the formation of a hydrogen bond with H of the aldehyde group in gossypol.

Rentgenografik tadqiqotlar izostrukturaviy klatratlarning ikkita guruhi mavjudligini aniqlab berdi, ular KK(A) va KK(V) kriptat guruhlariga tegishli deb aniqlandi. KK(A) kriptat guruhiga gossipolning benzol, brombenzol, izopropilbenzol, izopropilbrom va trixloretilen bilan hosil qilgan kriptatlar kiradi. Ushbu guruh kristallarida mezbun: mehmon nisbati 2:1, fazoviy guruhi R-1, z=4 ga teng.

Gossypol molecules are combined into dimers through hydrogen bonds, and later, these dimers are associated with columns using hydrogen bonds. When such columns are packed into the crystal structure, they form spaces where guest molecules are located.

Gossypol and its derivatives (mainly Schiff bases of gossypol) readily form complexes with many polar (only those with proton acceptor groups) and nonpolar substances. In addition, in clathration studies of gossypol derivatives, the phenomenon of polymorphism in the formation of host-guest complexes was revealed. It has been shown that gossypol derivatives can form different crystalline



modifications with the same guest component. The reason for this is that the formation of clathrate depends on the thermodynamic conditions of deposition.

Gossypol Schiff bases were reacted with Cu^{2+} , Ni^{2+} salts in a 1:2 mol ratio. The duration of the reaction was monitored hourly by thin-layer chromatography, and the obtained metallocomplexes were analyzed by comparison with the Schiff base spectra, and the following results were obtained. The main characterizing bond of the Schiff base is the azomethine bond formed between the carbonyl in gossypol and the primary amino group. When analyzing the IR-spectrum of the metallocomplex formed by Gossypol Schiff base with Cu^{2+} salt, it was concluded as follows. In contrast to the IR spectrum of the Schiff base, in the spectrum of the metallocomplex, it can be seen that the valence vibrations of the azomethine bond are shifted to the region of 1619 cm^{-1} . This displacement occurs when the nitrogen atom participating in the formation of an azomethine bond forms a donor-acceptor bond with Cu^{2+} in the metallocomplex.

It can also be seen that the region of 3485 cm^{-1} belonging to the Schiff-based $-\text{OH}$ group was shifted to the region of 3422 cm^{-1} in the spectrum of the metallocomplex, and a new absorption maximum appeared in the region of 3288 cm^{-1} . These changes are related to the formation of a covalent bond between the oxygen atom of 7-OH and Cu^{2+} on the Schiff basis. In the spectrum of the metallocomplex, we can consider the new absorption maxima at 544 cm^{-1} and 610 cm^{-1} as peaks belonging to metal-oxygen and metal-nitrogen bonds.

Table 2

Physicochemical quantities of Schiff base metallocomplexes

Synthesized metallocomplexes	solvent	d^{20}_S	Liquid refractive index	Color	Reaction product in (%)
Di-(2-amino-4-pikolin)gossipol + $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$	D MFA D MSO	291- 292	,49	brown	67 .8
Di-(2-amino-4-pikolin)gossipol + $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$	D MFA D MSO	297- 298	,44	orange yellow ashil	69 .3
Di-(6-amino-3-pikolin)gossipol + $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$	D MFA D MSO	300<	,54	brown	71 .3
Di-(6-amino-3-pikolin)gossipol + $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$	D MFA D MSO	300<	,69	brown	70 .8

System:- hexane:acetone (4:1)

Of the Schiff base formed by gossypol with 6-amino-3-picoline When studying the UV spectrum, 236,51 for this substance; It gave absorption maxima at 379,1 nm. When analyzing the UV-spectrum of



the metallocomplex of the Schiff base of Gossypol with 6-amino-3-picoline obtained with $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$, the metallocomplex is 247,64; It gave absorption maxima at 422,81 nm. The graph comparing these spectra to each other looked like this:

236.51 based on Schiff; The peak of the absorption maxima at 379.1 nm is lost in the metallocomplex, and the Schiff-based peak at 379.1 nm is shifted to 422.81 nm in the metallocomplex, this shift occurs due to coordination bonds in the metallocomplex.

Of the Schiff base formed by gossypol with 6-amino-3-picoline When analyzing the IR spectrum, obvious changes occur in the area of absorption maxima belonging to the NH_2 group at $3120,44 \text{ cm}^{-1}$. Based on the Schiff, we can see that the surface and distance between these peaks has expanded. These expansions occur due to Schiff-based hydrogen bonds. As a result, we can see absorption maxima at $1714,68 \text{ cm}^{-1}$ due to valence vibrations of the new $-\text{N}=\text{CN}-$ bond.

It is a Schiff base and a metallocomplex formed with $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$ When the IR spectrum was analyzed, it was found that the absorption maxima at $2966,25 \text{ cm}^{-1}$ due to valence vibrations of the new $-\text{N}=\text{CN}-$ bond did not change and $3479,34$; Absorption maxima were observed at $3411,77 \text{ cm}^{-1}$ due to coordination bonds in the metallocomplex. In the IR-spectra of the obtained metallocomplexes, the change of the valence vibration lines belonging to the OH group (area $2740\text{-}3300 \text{ cm}^{-1}$) and the weakening of the vibration frequencies of azomethine bonds (area $1600\text{-}1627 \text{ cm}^{-1}$) is observed; M-O (area $450\text{-}490 \text{ cm}^{-1}$) lines appear.

When studying the structure of metal complexes using UV spectroscopy, the presence of azomethine bonds ($\lambda_{\text{C}=\text{N}}=272 \text{ nm}$) is proven by the appearance of an absorption line in the 260-300 nm region.

Literature

1. Hakberdiev, S. M., Talipov, S. A., Dalimov, D. N., & Ibragimov, B. T. (2013). 2, 2'-Bis {8-[(benzylamino) methylidene]-1, 6-dihydroxy-5-isopropyl-3-methylnaphthalen-7 (8H)-one}. *Acta Crystallographica Section E: Structure Reports Online*, 69(11), o1626-o1627.
2. Хакбердиев Ш. М., Тошов Х. С. Моделирование реакции конденсации госсипола с ортолуидином //ББК 74.58 G 54. – С. 257.
3. Khamza, Toshov, Khakberdiev Shukhrat, and Khaitbaev Alisher. "X-ray structural analysis of gossypol derivatives." *Journal of Critical Reviews* 7.11 (2020): 460-463.
4. Хакбердиев Ш. М., Асророва З. С. Ғўза илдизидан госсипол олиш, госсипол ҳосилалари синтези ва тузилиши //Science and Education. – 2020. – Т. 1. – №. 2.
5. Хакбердиев, Ш. М. (2020). Бензиаминнинг госсиполли ҳосиласи синтези, тузилиши ва мис, никель, собаль тузлари билан металлокомплексларини олиш. *Science and Education*, 1(8), 16-21.
6. Хакбердиев, Ш. М., & Муллажонов, З. С. Қ. (2020). Госсипол ҳосилаларининг паренхиматоз аъзолар тўқималари ва макрофаглар миқдорига таъсири. *Science and Education*, 1(9).
7. Хакбердиев, Ш. М. (2020). Турли тузилишли аминларнинг госсиполи ҳосилалари синтези ва биологик фаоллиги. *Science and Education*, 1(9).
8. Khakberdiyev, S. M. (2021). Study of the structure of supramolecular complexes of azomethine derivatives of gossypol. *Science and Education*, 2(1), 98-102.
9. Ҳамидов С. Х., Муллажонов З. С. Қ., Хакбердиев Ш. М. Кумушнинг госсиполли комплекси ва спектрал таҳлили //Science and Education. – 2021. – Т. 2. – №. 2.



10. Хақбердиев Ш. Янги шифф асослари ва уларнинг сувда эрувчан комплекслари тузилишини ўрганиш //Журнал естественных наук. – 2021. – Т. 1. – №. 2.
11. Хамидов, С. Х., & Хақбердиев, Ш. М. (2021). Бирламчи алифатик аминларнинг госсиполли ҳосилалари синтези. *Science and Education*, 2(3), 113-118.
12. Муллажоновна, З. С., Хамидов, С. Х., & Хақбердиев, Ш. М. (2021). Турли усулларлар ёрдамида госсиполли комплекс таркибидан кумуш ионини аниқлаш. *Science and Education*, 2(3), 64-70.
13. Khaitbaev A. K., Khakberdiev S. M., Toshov K. S. Isolation of Gossypol from the Bark of Cotton Roots //Annals of the Romanian Society for Cell Biology. – 2021. – С. 1069-1073.
14. Хақбердиев Ш. Госсипол ҳосилалари, металлокомплекслари синтези қилиш ва кукунли дифрактометрда ўрганиш //Журнал естественных наук. – 2021. – Т. 1. – №. 2.
15. Хақбердиев Ш. Шифф асоси ва металлокомплексларининг термик анализи //Журнал естественных наук. – 2021. – Т. 1. – №. 3.
16. Хақбердиев Ш. Синтез, строение и получение супрамолекулярных комплексов ароматических аминов с госсиполом //Журнал естественных наук. – 2021. – Т. 1. – №. 4.
17. Хақбердиев Ш. М. и др. Синтез госсипольных производных орто, мета, пара толуидина и их строение //Science and Education. – 2021. – Т. 2. – №. 10. – С. 195-200.
18. Khakberdiev, Sh M., et al. "Synthesis and structure of gossypol azomethine derivatives." *Young Scientist*,(4) (2015): 42-44.
19. Хақбердиев Ш. М. и др. 3-аминопропанол-1 билан госсиполнинг турли комплекслари синтези ва макрофаглар микдориға таъсири //Журнал естественных наук. – 2021. – Т. 1. – №. 1.
20. Хақбердиев, Ш. М. (2021). Госсиполнинг аминопиридинлар билан синтези ва уларнинг никел тузи металлокомплексларини олиш. *Журнал естественных наук*, 3(5), 10-15.
21. Хақбердиев, Ш., Қодир, Д., Маматова, Ф., & Муллажоновна, З. (2022). Госсипол асосида ациклик аминобирикмаларнинг ҳосилалари синтези. *Журнал естественных наук*, 1(2 (7)), 12-16.
22. Mahramovich, K. S., Sattarovna, K. F., & Farangiz, M. (2022). Synthesis of Gossipy Products of Pyrimidine Bases and Getting Their Water-Solved Complexes. *Eurasian Scientific Herald*, 8, 118-121.
23. Mahramovich, K. S. (2022). Results of computer study of biological activity of gossipol products. *Web of Scientist: International Scientific Research Journal*, 3(6), 1373-1378.
24. Хақбердиев, Ш., Муллажоновна, З., & Маматова, Ф. (2022). Адениннинг госсиполли ҳосиласи унинг металл ва супрамолекуляр комплексларини турли таҳлиллар асосида ўрганиш. *Журнал естественных наук*, 1(2 (7)), 288-293.
25. Khakberdiyev Shukhrat Mahramovich, & Mamatova Farangiz Qodir qizi. (2022). Synthesis of metallocomplexes of schiff bases and their structural analysis. *World Bulletin of Public Health*, 16, 173-177. Retrieved from
26. Mahramovich, K. S. (2023). Structural analysis of supramolecular complexes of schiff bases. *American Journal of Interdisciplinary Research and Development*, 12, 36-41.
27. Khakberdiyev Shukhrat Mahramovich, Azizova Safina Isroiljon qizi, Mamatova Farangiz Qodir qizi, Rabbimova Marjona Ulug'bek qizi. (2023). Biological Activities of Water-Soluble and Cu²⁺ Salts of Gossypol Derivatives Metallocomplexes. *International Journal of Scientific Trends*, 2(2), 55–60. Retrieved from



28. Mahramovich, K. S., & Khodiyevich, K. S. (2023). Study of the practical significance of benzimidazole and some of its derivatives. *Open Access Repository*, 4(02), 80-85.
29. Ramírez-Coronel, A. A., Mezan, S. O., Patra, I., Sivaraman, R., Riadi, Y., Khakberdiev, S., ... & Fakri Mustafa, Y. (2022). A green chemistry approach for oxidation of alcohols using novel bioactive cobalt composite immobilized on polysulfone fibrous network nanoparticles as a catalyst. *Frontiers in Chemistry*, 10, 1015515.
30. Khakberdiev Shukhrat Mahramovich, Khamidov Sobir Khodiyevich. (2023). Chemical structure and practical significance of benzoxazole . *Ethiopian International Journal of Multidisciplinary Research*, 10(09), 75–77.
31. Mahramovich, K. S. (2024). Study of synthesis, structure and biological activity of gossypol derivatives in computer program. *American Journal of Innovation in Science Research and Development*, 1(2), 75-81.
32. Муллажонова, З. С. Қ., Хамидов, С. Х., & Хакбердиев, Ш. М. (2021). Турли усулларлар ёрдамида госсиполли комплекс таркибидан кумуш ионини аниқлаш. *Science and Education*, 2(3), 64-70.

