



Global Longitudinal Strain is an Informative Index of Left Ventricular Performance in Neonates Receiving Intensive Care

Shodiyev Mirshodbek Marufovich

Bukhara State Medical Institute, Assistant of the Department of Internal Medicine in Family Medicine. Bukhara, Republic of Uzbekistan

Abstract: Echocardiographic assessment of left ventricular function is crucial in NICU. The study aimed to compare the accuracy and agreement of global longitudinal strain (GLS) with conventional measurements. Real-life echocardiograms of neonates receiving intensive care were retrospectively reviewed. Shortening fraction (SF), ejection fraction (EF) and S' measurements were retrieved from health records. GLS was calculated offline from stored images. The association with stroke volume indexed for body weight (iSV) was evaluated by regression analysis.

Key words: Ventricular Function, Left, Echocardiography, Global Longitudinal Strain, Intensive care, neonatal, Infant, Newborn.

This was a single-center retrospective study conducted at a tertiary referral academic neonatal intensive care unit (NICU). Ethics approval was obtained from the local Research Ethics Committee. Parents or guardians of neonates gave their written informed consent. This was a convenience sample: all neonates, admitted from

January to December 2022, who underwent echocardiographic assessment of LV function were included in the study. Exclusion criteria were: major congenital abnormalities, and congenital heart disease, except for patent ductus arteriosus (PDA) and patent foramen ovale. During the study period, NPE was used to assess neonates with PDA, persistent pulmonary hypertension of the newborn (PPHN), shock and/or hypoperfusion, for the screening of congenital heart disease (CHD) and pulmonary hypertension in neonates with bronchopulmonary dysplasia (BPD) and research purposes. All examinations were performed using a Philips Epiq 7 system with 12 MHz probe (Philips Ultrasound, Andover, MA, USA). Images were acquired according to the recommendations of the American Society of - Echocardiography (13) .

Electrocardiograms were recorded simultaneously. Three cardiac cycles were digitally stored at high frame rate (100–130 Hz). Image quality was assessed according to Colan et al.(14) Images considered poor or unusable were excluded from the - analysis (15) .

Conventional measurements of LV systolic function. Conventional measurements of LV systolic function were retrieved from electronic health records. Measurements were done offline, immediately after scanning, using the QLAB software available on the ultrasound system. The following measurements were obtained as previously described: SF was calculated from M-mode images of a short axis view at the level just distal to the leaflet tips of the mitral valve (Fig. 1a) TDI was used to measure myocardial velocities from an apical 4-chamber view, placing the sample volume just apical to the lateral margin of the mitral valve: systolic peak velocity (S') was measured (Fig. 1b) EF was calculated from a 4- and 2-chamber view based on the modified biplane Simpson's method of disc (Fig. 1c)

Global Longitudinal Strain (GLS)

According to our echocardiographic protocol, an apical 4-, 3- and 2-chamber view with ECG trace was recorded, trying to obtain an optimal visualization of the LV wall and to avoid foreshortening of the LV. The analysis of digitally stored images was performed offline using the Philips QLAB software available on the ultrasound system, version 10.0 (https://www.usa.philips.com/healthcare/products/HCNOC_TN145/qlab-cardiovascular-ultrasound-quantification-software-documents), by an operator blind to the conventional echo data, at least 4 weeks apart from image acquisition. As previously described, the QRS complex was used as a reference time-point. The region of interest (ROI), including six segments with different colors, was automatically traced by an endocardial point-and-click approach in each apical view (4-, 3- and 2-chamber view). The ROI was manually corrected, if needed, Then the software automatically tracked the movement of speckles to derive the deformation parameters. The 18-segment model was used; peak longitudinal strain from a 4-, 3- and 2-chamber view and GLS were calculated as previously reported (Fig. 1d)16.

Stroke volume

Stroke volume (SV) was used to assess the accuracy of the measurements of LV systolic function as previously described in - adults17. To calculate SV, aortic internal diameter was measured at the valve hinge points from a parasternal long axis view and aortic velocity time integral was measured from an apical 5-chamber view. SV was indexed for body weight (iSV)(18,19).

Statistical analysis

The neonates enrolled in the study were divided into two subgroups: neonates with impaired and normal LV function. The latter was defined as $GLS \leq -15$ 10,20. Normality of data distribution was assessed by Shapiro–Wilk’s test. Data are presented as mean (\pm SD), median (interquartile range (IQR)) or count (%) for parametric, non-parametric and categorical variables, respectively. Differences between the two groups of neonates were evaluated using an independent-sample t-test, Mann–Whitney U test and χ^2 test in case of parametric, non-parametric, and categorical variables, respectively. Pearson’s partial correlation was run to assess the relationship between SF, EF, S’, GLS and iSV. Univariate and multivariate linear regression analysis was run to estimate the ability of the conventional LV systolic measurements (SF, EF, S’) and that of GLS to predict the iSV. Receiver-operating characteristic (ROC) curve analysis was used to assess the diagnostic ability of GLS, EF, S’ and SF to identify neonates with uncompensated shock. The latter was defined as blood pressure below the 3rd centile according to gestational age and clinical signs of hypoperfusion (prolonged capillary refill time, tachycardia, tachypnea, mottled skin and/or cool extremities, reduced or absent urine output, lactic acidosis)(21). Cohen’s κ was run to assess the agreement between GLS and the other conventional measurements (SF, EF and S’) on whether the left ventricular systolic function was judged impaired or normal. According to our institutional echocardiographic protocol, the following cut-off values to define impaired LV function were used as previously described: $SF \leq 25\%$, $EF \leq 55\%$, $S' \leq 3$ cm/s in neonates with gestational age (GA) < 30 weeks, $S' \leq 3.5$ cm/s in neonates with GA from 30 to 36 weeks, $S' \leq 4.2$ cm/s in neonates with GA ≥ 37 weeks. For the advanced echocardiographic imaging, we used a $GLS > -15$ to define an impaired LV - function5,16,2

Figure 1. (a) M-mode image of a short axis view at the level just distal to the leaflet tips of the mitral valve showing shortening fraction (SF) calculation. (b) Tissue doppler imaging of an apical 4-chamber view with the sample volume at the lateral margin of the mitral valve, showing systolic peak velocity (S’) (c) Calculation of ejection fraction (EF) according to Simpson’s biplane method of disc. Apical 4-chamber views (top) showing the measurement of left ventricular end-diastolic (left) and end-systolic (right) volumes. Apical 2-chamber view (bottom) showing the measurement of left ventricular end-diastolic (left) and end-systolic volumes (right). (d) Longitudinal-Strain in 2-chamber view (top-left), 3-chamber view (top-right) and 4-chamber view (bottom-left). Bull’s-eye of global



longitudinal strain (GLS). Bull's eye plot derived from the longitudinal strain of the three apical views (bottom-right).

Ethics approval

The study was approved by the Research Ethics committee of the University Hospital of Verona, no. 2132CESC. The parents or guardians of babies enrolled gave their written informed consent. All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/ or national research committee and with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards.

Results

From January to December 2022 a total of 460 echocardiograms of 215 neonates were performed and were assessed for eligibility. One hundred and ten echocardiograms of 44 neonates did not meet the inclusion criteria: in 39 echocardiograms a complete assessment of LV function was not performed, in 26 echocardiograms no stored images were found, lastly, 33 and 12 echocardiograms were excluded due to CHD and major congenital abnormalities, respectively. Following the image quality assessment of the 4-, 3- and 2-chamber views of stored clips, 16 echocardiograms of 16 neonates were excluded. Finally, 334 echocardiograms of 155 neonates were included in the analysis. The study flow chart is shown in Supplemental Fig. 1. The echocardiograms were performed for the following reasons: PDA assessment 108 (30.9%), screening CHD 101 (28.9%), screening BPD 39 (11.1%), shock and/or hypoperfusion 20 (5.7%); research purposes 78 (22.3%), PPHN 4 (1.1%).

	Impaired LV function (72 neonates)	Normal LV function (262 neonates)	P value
Gestational age (week)	34.4 ± 4.7	34.9 ± 4.1	0.36
Weight at birth (g)	2357 ± 1114	2289 ± 899	0.59
Weight at scan (g)	2348 ± 1101	2301 ± 858	0.70
Female	31 (43.1%)	113 (43.5%)	.95
Day of life	2 [1-2]	1 [1-2]	0.02
Ventilation			
None	38 (52.8%)	148 (56.5%)	0.66
Non-invasive ventilation	19 (26.4%)	102 (38.9%)	0.06
Invasive ventilation	15 (20.8%)	12 (4.6%)	<0.01
FiO ₂	0.21 [0.21-0.23]	0.21[0.21-0.21]	0.67
iNO	9 (12.5%)	6 (2.3%)	0.01
Systolic blood pressure (mmHg)	66 ± 14	68 ± 9	0.52
Mean blood pressure (mmHg)	45 ± 11	47 ± 8	0.22
Diastolic blood pressure (mmHg)	37 ± 11	39 ± 9	0.44
Daily total fluid (ml/kg/die)	103 ± 42	94 ± 40	0.13
Vasopressor and/or inotrope	10 (13.9%)	4 (1.5%)	<0.01
Sepsis at the time of echo	12 (16.7%)	25 (9.5%)	0.09
IUGR	4 (5.6%)	33 (12.6%)	0.13
Antenatal steroids	21 (29.2%)	53 (20.2%)	0.11

Table 1. Clinical characteristics at the time of echo of the two cohorts of neonates with normal and impaired GLS. Values are presented as mean ± SD, median [IQR] and count (%). LV left ventricle, iNO inhaled nitric oxide, IUGR intrauterine growth restriction. Significant values are in bold.

Discussion

Monitoring the adequacy of the oxygen delivery to cells and tissues and applying the appropriate therapeutic interventions is a dogma of intensive care medicine. Longitudinal monitoring of SV, which is the amount of blood ejected through the aortic valve during systole, is of paramount importance. SV, together with heart rate and arterial oxygen content, is a determinant of oxygen delivery 3. As the summative event of the cardiac cycle, from the perspective of the critical care physician, SV is the most pragmatic indicator of the well-being of the heart and the cardiovascular system, independent of - EF19,23 . In this study, the association with iSV of GLS and the other conventional measurements of LV function, their ability to identify neonates with uncompensated shock, and their agreement were evaluated for the first time in real-life echocardiograms of neonates requiring intensive care. Firstly, a strong association of iSV with GLS was found, in contrast to the weaker associations with SF, EF and S'. Secondly, when GLS and all the conventional parameters



were included in the regression model to predict iSV, more than 70% of the variance of iSV was explained by GLS, with only a limited contribution from S', EF and SF. Thirdly, the ability of GLS to identify neonates with uncompensated shock was superior compared to the other measurements of LV function. Finally, a limited agreement was found between GLS and the conventional echocardiographic measures of systolic function. Our data confirmed that SF, EF and S' are far from being accurate in the neonatal population, showing only a moderate correlation with iSV and a limited ability to identify neonates with uncompensated shock. SF was the measure that showed the lowest accuracy. It relies on normal LV septal morphology, the latter being commonly altered in neonates, due to the increased pulmonary pressure after birth. Moreover, previous findings showed that the longitudinal shortening of the LV is the major contributor to the - SV17. SF, which assesses the radial shortening, is of little value. We feel that its use, as a single parameter of LV function, should be discouraged in neonates, although the available guidelines still include SF in the quantitative assessment of LV - function 2,3 . The accuracy of both S' and EF was slightly superior but still suboptimal. S' evaluates the myocardial velocities during the cardiac cycle only in one specific region of the heart, adjacent to the mitral annulus, it is angle- dependent and cannot differentiate active from passive - motion5 . EF calculation with Simpson's biplane method of disc is based on geometric assumptions, which may lead to inaccuracies in measurements, especially in dilated ventricles or in the presence of abnormal septal - motion (24) . Although EF is a cornerstone in the assessment of LV systolic function and the management of heart failure, there is growing evidence suggesting that EF is not a sensitive marker of myocardial dysfunction. EF, which measures changes in the volume of the LV cavity, is only an indirect measure of LV function, and is prone to geometrical confounders, specifically in the case of LV hypertrophy or reduced LV - cavity (25). Our data underlines that, from the neonatologist's perspective, there is an urgent need for more accurate measures of LV systolic function, due to the poor accuracy of the conventional measures and their limited agreement with GLS. We believe that advanced echocardiographic imaging is a promising tool. GLS provides a direct measure of myocardial systolic - function (26). We acknowledge that the inter-vendor variations and the need for off-line analysis can inhibit the adoption of GLS in clinical practice. However, future technical developments and the availability of automated analysis may facilitate this process. In the current study, GLS was calculated using the software available on the ultrasound system, as in the real-world assessment of LV function at the cotside, showing an optimal potential for the clinical application of this technique. GLS showed a strong association with iSV at the univariable analysis and was the major determinant of the variance of iSV, compared to the conventional measurements, at the multivariate analysis. We hypothesized that GLS is a more sensitive index of LV dysfunction compared to EF due to the complex myocardial architecture with three distinct layers of fibers with different - orientations (27). GLS allows a better assessment of the deformation of the longitudinal sub-endocardial fibers, which are more vulnerable to wall stress and mostly contribute to stroke - volume(17). Moreover, it overcomes the shortcomings of the conventional measurements: it does not rely on geometrical assumption, it is not angle-dependent, it differentiates between active and passive motion of the heart, and it detects impaired myocardial function even in the presence of LV hypertrophy and reduced LV - volume (25). In adults and older children, there is growing evidence of the prognostic and diagnostic superiority of GLS over conventional parameters in a wide range of diseases, with optimal reproducibility. Previous data confirmed that myocardial deformation imaging is feasible, reproducible and accurate also in neonates with PDA, PPHN, hypoxic ischemic encephalopathy, and congenital diaphragmatic - hernia16,28-31. Our findings provide evidence that GLS should be added to the armamentarium of the NPE.

Conclusions. Using real-life echocardiograms from a large cohort of neonates requiring intensive care we showed that GLS was a more informative index of LV performance compared to the conventional measures (EF, SF, S'). Our findings provide the pathophysiological and clinical rationale for a more extensive use of GLS in the quantitative assessment of LV systolic function, to guide management. Further studies are needed to evaluate the association with relevant outcomes and



the efficacy of therapeutic interventions, before widespread incorporation of GLS into mainstream clinical practice.

References

1. Corsini, I. et al. Neonatologist performed echocardiography (NPE) in Italian neonatal intensive care units: A national survey. *Ital. J. Pediatr.* 45, 131. <https://doi.org/10.1186/s13052-019-0721-z> (2019).
2. Mertens, L. et al. Targeted Neonatal Echocardiography in the Neonatal Intensive Care Unit: Practice Guidelines and Recommendations for Training: Writing group of the American Society of Echocardiography (ASE) in collaboration with the European Association of Echocardiography. *Eur. J. Echocardiogr.* 12, 715–736. <https://doi.org/10.1093/ejechocard/je181> (2011).
3. de Boode, W. P. et al. The role of neonatologist performed echocardiography in the assessment and management of neonatal shock. *Pediatr. Res.* 84, 57–67. <https://doi.org/10.1038/s41390-018-0081-1> (2018).
4. Singh, Y. et al. International evidence-based guidelines on Point of Care Ultrasound (POCUS) for critically ill neonates and children issued by the POCUS working group of the European society of paediatric and neonatal intensive care (ESPNIC). *Crit. Care* 24,65. <https://doi.org/10.1186/s13054-020-2787-9> (2020).
5. Nestaas, E., Schubert, U., de Boode, W. P. & EL-Khuffash, A. Tissue Doppler velocity imaging and event timings in neonates: A guide to image acquisition, measurement, interpretation, and reference values. *Pediatr. Res.* 84, 18–29. <https://doi.org/10.1038/s41390-018-0079-8> (2018).
6. Leo Lopez, M. D. F. C. et al. Recommendations for quantification methods during the performance of a pediatric echocardiogram: A report from the pediatric measurements writing group of the American society of echocardiography pediatric and congenital heart disease council. *J. Am. Soc. Echocardiogr.* 23, 465–495. <https://doi.org/10.1016/j.echo.2010.03.019> (2010).
7. Tissot, C., Singh, Y. & Sekarski, N. Echocardiographic evaluation of ventricular function—For the neonatologist and pediatric intensivist. *Front. Pediatr.* <https://doi.org/10.3389/fped.2018.00079> (2018).
8. Klaeboe, L. G. & Edvardsen, T. Echocardiographic assessment of left ventricular systolic function. *J. Echocardiogr.* 17, 10–16. <https://doi.org/10.1007/s12574-018-0405-5> (2019).
9. JainEL-Khuffash, A. A. F. et al. Left ventricular function in healthy term neonates during the transitional period. *J. Pediatr.* 182, 197–203.e2. <https://doi.org/10.1016/j.jpeds.2016.11.003> (2017).
10. LevyEL-Khuffash, P. T. A. et al. Maturation patterns of systolic ventricular deformation mechanics by two-dimensional speckle-tracking echocardiography in preterm infants over the first year of age. *J. Am. Soc. Echocardiogr.* 30, 685–698. <https://doi.org/10.1016/j.echo.2017.03.003> (2017).
11. Breatnach, C. R. et al. Novel echocardiography methods in the functional assessment of the newborn heart. *Neonatology* 110, 248–260. <https://doi.org/10.1159/000445779> (2016).
12. Ficial, B. et al. Tissue-tracking mitral annular displacement in neonates. *J. Ultrasound Med.* <https://doi.org/10.1002/jum.16399> (2023).
13. Lai, W. W. et al. Guidelines and standards for performance of a pediatric echocardiogram: A report from the task force of the pediatric council of the American society of echocardiography. *J. Am. Soc. Echocardiogr.* 19, 1413–1430. <https://doi.org/10.1016/j.echo.2006.09.001> (2006).



14. Colan, S. D. et al. The ventricular volume variability study of the pediatric heart network: Study design and impact of beat averaging and variable type on the reproducibility of echocardiographic measurements in children with chronic dilated cardiomyopathy. *J. Am. Soc. Echocardiogr.* <https://doi.org/10.1016/j.echo.2012.05.004> (2012).
15. Ficial, B. et al. Feasibility, reproducibility and reference ranges of left atrial strain in preterm and term neonates in the first 48 h of Life. *Diagnostics* <https://doi.org/10.3390/DIAGNOSTICS12020350> (2022).
16. EL-Khuffash, A. et al. Deformation imaging and rotational mechanics in neonates: A guide to image acquisition, measurement, interpretation, and reference values. *Pediatr. Res.* 84, 30–45 (2018).