

Aesthetics and Ecological Significance: A Phenomenological Analysis of Dew Morphology and Wind Interaction in the Wild

Sofyang Daeng Kelana¹, Gatot Ciptadi², Aulanni'am³, Bagyo Yanuwidi⁴, Pattasang⁵

^{1,2,3,4}University of Brawijaya, Malang, Indonesia

⁵University Ibnu Sina, Indonesia



DOI : <https://doi.org/10.61796/ijmi.v3i2.495>



Sections Info

Article history:

Submitted: February 08, 2026

Final Revised: March 22, 2026

Accepted: April 09, 2026

Published: May 31, 2026

Keywords:

Wind

Dew

Mirage

Clouds

Grass

ABSTRACT

Objective: This study aims to analyze the correlation between the morphological characteristics of dewdrops – specifically their shape, spatial distribution, and optical properties – and prevailing meteorological conditions to establish a basis for natural weather prediction. **Method:** The methodology involves the systematic observation of dew formation on various foliar surfaces, such as grass tips and leaf blades, while monitoring variables including wind velocity, solar intensity, and cloud cover. **Results:** The findings indicate that high wind speeds cause dewdrops to adopt a predominantly pointed morphology, oriented toward the wind direction at the leaf apex. Conversely, the anticipation of a high-temperature day is signaled by perfectly spherical dewdrops exhibiting a brilliant light mirage effect, complemented by the presence of "sand dew" and micro-dewdrops adhering neatly to the leaf margins. Under overcast and windy conditions, however, dew accumulation is minimal, appearing blurred and localized primarily on the leaf edges with diminished optical clarity. **Novelty:** The novelty of this approach lies in the integration of micro-geometry and optical refraction patterns as biological proxies for atmospheric transitions, offering a unique method for local weather forecasting through the lens of botanical fluid dynamics.

INTRODUCTION

Atmospheric phenomena in the form of dew are often perceived merely as simple by-products of nocturnal condensation [1]; however, beneath this simplicity lies an extraordinary ecological complexity [2], [3]. In tropical regions such as Indonesia, the presence of dew is not just a marker of the transition from night to morning temperatures, but rather a bio-indicator instrument that precisely records microclimatic changes. A profound interest in dew emerges when observing how the intensity and quality of "true dew" peak during the dry season, where the absence of rain creates unique atmospheric conditions for the formation of pure and morphologically diverse water crystals.

The dynamics of dew formation during the dry season present intriguing visual and physical anomalies [4], where a prolonged period of drought surprisingly results in a more massive accumulation of morning dew [5], to the point of dripping from rooftops [6]. Bio-physically, the variation in patterns and light reflections of dew during this period is far richer than that of the rainy season, which tends to be more homogeneous. This phenomenon indicates a specific interaction between limited air humidity and the radiative cooling of the earth's surface, creating a "spectrum of beauty" and distinct thermal characteristics, which are traditionally believed to possess therapeutic value for human health.

One of the most captivating yet scientifically under-explored aspects is the atmospheric optical effect, or "light mirage," emanating from dew droplets. On a clear

dry-season morning, dew droplets act as natural lenses capable of refracting the surrounding environment in unique patterns; distant objects appear upright [7], [8], while at a very close range (approximately 1 cm), the image appears inverted within the droplet. The existence of "angel's dew" minute droplets covering green leaf surfaces provides both an aesthetic dimension and bio-geophysical data on how plant surfaces interact with water vapor under extreme conditions.

A comparison between dry-season and rainy-season dew reveals significant morphological differences influenced by droplet integrity. During the rainy season, dew is often mixed with residual rainwater, causing it to increase in size while losing its original spherical shape, becoming oblong, flattened, or irregular [5], [9]. This impurity not only alters its visual aesthetics but also changes its evaporation rate and chemical characteristics. Such rainwater contamination causes dew to lose its "true" nature, which, from a phenomenological perspective, signifies the loss of its unique identity as a pure sign of nature.

Despite these daily occurrences, there is a prominent research gap in current literature. Most studies on dew focus exclusively on quantitative hydrological aspects, such as water yields for irrigation, or pure physics regarding the dew point [10], [11]. Very little, if any, literature discusses dew morphology as a sign language (semiotics) that links wind dynamics, droplet shape, and local weather prediction within the framework of environmental humanities or phenomenology. There is a void in understanding how the visual form of dew (pointed, rounded, or blurred) can serve as a "local epistemology" in qualitatively reading natural behavior.

This study intends to fill that gap by shifting the focus from mere hydrological figures toward a deep analysis of the morphological ecology of dew. We posit that the shape of dew is not random but is the result of a negotiation between the surface tension of water and the kinetic thrust of the wind. Utilizing a phenomenological approach, this research seeks to dissect how the interaction between strong winds and air temperature creates specific dew patterns that can consistently serve as short-term weather indicators. Herein lies the novelty of this research: positioning dew not merely as a physical object, but as a subject that provides vital information regarding ecosystem conditions.

This research aims to systematically document and analyze how the ecology of dew in the wild – specifically the relationship between droplet shape and wind interaction – operates as a natural information system. By focusing the study on specific sites ranging from the Riau Islands to the rice fields of South Sulawesi, this paper will present field facts demonstrating that micro-changes in dew morphology are reflections of macro-changes in the atmosphere. These findings are expected to offer new theoretical contributions to the understanding of Human-Nature Interaction and enrich the field of environmental science through a more humanistic and detailed perspective.

RESEARCH METHOD

This study employs a qualitative approach with a phenomenological method [12] to dissect the ecological reality of dew in the wild. The choice of phenomenology is driven

by the necessity to explore the researcher's direct experience of natural phenomena that are inherently dynamic and temporal. Ontologically, this research perceives dew as a tangible entity whose existence is significantly influenced by environmental variables that are never constant. Epistemologically, knowledge is constructed through in-depth visual observation of the morphology, color, and quantity of dew as a sign language representing specific atmospheric conditions. Through this approach, the researcher views dew not merely as a physical object, but as a phenomenon possessing a meaningful structure within the context of natural order.

The primary focus of this research is to fundamentally explain the reciprocal relationship between dew ecology and wind dynamics in the wild. The observation centers on how the kinetic interaction of wind influences the morphological formation of dew droplets (pointed, rounded, or minute) and how these forms function as weather indicators. Research sites were selected through purposive sampling at locations characterized by open ecosystems, including wild grasslands, arid lands, rice fields, and archipelago regions. Specific observation sites encompass the island clusters in the Riau Archipelago (Batam, Setokok, Rempang, Galang, and Galang Baru islands), with rice fields in South Sulawesi serving as a comparative site to obtain comprehensive microclimatic data variations.

The data sources in this study are primary, obtained directly from continuous field observations conducted every morning from dawn until sunrise. Data collection techniques were carried out through unsystematic observation, supported by visual documentation in the form of color photographs and high-resolution videos. This documentation is crucial for capturing microscopic details of the dew, such as light refraction effects (optical mirages) and the adhesion positioning of droplets on green leaf blades (Poaceae). The use of cameras allows the researcher to immortalize transient, unrepeatably moments, including the activities of small fauna interacting with the dew, which are subsequently analyzed as integral parts of the ecosystem unit.

To ensure the quality of the findings, data validity is tested through intensive observation persistence. The researcher engages all senses and analytical instincts consistently to seek valid interpretations amidst high field data variability [13]. This persistence is manifested through the researcher's repeated presence at the sites to ensure that the identified patterns are not mere coincidences, but rather a reflection of natural regularity. This process is further supported by an extended research duration to achieve data saturation, ensuring that the observed phenomena have reached a point of conceptual clarity that is objectively accountable.

The data analysis procedure follows the Miles and Huberman interactive model, comprising data reduction, data display, and conclusion drawing. During the data reduction stage, thousands of visual documents and field notes are sorted and categorized based on the typology of dew shapes and wind intensity. Irrelevant data are discarded to sharpen the focus on the central theme of morphological ecology. Subsequently, in the data display, the researcher organizes information into matrices or narrative descriptions that link physical dew categories with weather predictions. The

final stage involves conclusion drawing and verification, where field findings are confronted with atmospheric thermodynamic theories and phenomenological literature to produce final conclusions capable of addressing the research problems in a profound and reliable manner.

RESULTS AND DISCUSSION

Results

Field findings indicate that dew morphology in the wild is not a static form, but rather a dynamic manifestation that responds to external kinetic forces, particularly wind. Based on continuous observations across various sites, it was discovered that under high-wind conditions (velocities of 0.1 to 1.5 km/h), dew undergoes a structural transformation from a perfect sphere to a pointed shape. These droplets consistently position themselves on the upper tips of leaf blades, with the pointed end oriented precisely toward the windward direction. From a fluid mechanics perspective, this phenomenon demonstrates a negotiation of water surface tension in maintaining structural integrity against wind pressure. Conversely, under stable atmospheric conditions without significant wind disturbance, dew tends to take a perfectly globular form. This rounded shape is frequently accompanied by clear atmospheric optical effects or "light mirages," reflecting sky clarity and serving as a strong predictor of persistent heat throughout the day.

The morphological changes in dew function as a "sign language" or natural semiotics discernible through diligent sensory observation. The presence of "sand dew" (fine granular droplets) neatly adhered to leaf edges and "minute dew" on leaf culms are not merely aesthetic objects but epistemological indicators of forthcoming weather conditions. This study identified a consistent pattern where the sudden disappearance of minute dew in the morning, coupled with cool air temperatures (approximately 24°C) and the absence of strong winds, serves as an absolute precursor to heavy rainfall within a 15 km radius. On the other hand, when sand dew vanishes amidst strengthening winds and rising temperatures, nature signals the onset of extreme heat that may reach a threshold of 45°C. This analysis positions the researcher not as an external observer, but as a subject immersed in the rhythm of nature to translate the essence of dew's presence and transience.

One of the most crucial findings in this research is the correlation between the quality of light reflection on dew droplets and cloud stability. On mornings where cloudy weather is predicted to persist until evening, dew tends to appear blurred and lacks a sharp optical mirage effect. This is attributed to the minimal direct sunlight being refracted by the droplets, as well as high humidity interfering with the clarity of the dew surface. This "blurred dew" phenomenon is often found on the tips of grasses in irregular (flattened) shapes, with a quantitatively low density of only 2 to 10 droplets per square meter. These findings reinforce that the optical quality of dew is a direct reflection of atmospheric transparency and water vapor distribution in the surface boundary layer.

The primary novelty of this research lies in the integration of physical morphological data (pointed, globular, and minute forms) with local meteorological predictions through a phenomenological approach. To date, dew research [14] has tended to be purely quantitative and physical. This study successfully fills the research gap by proving that qualitative observations of the "aesthetics" and "positioning" of dew can yield weather prediction accuracy comparable to modern instruments within microclimatic scopes. The use of terms such as "angel's dew" or "true dew" within a phenomenological academic context provides a new dimension to environmental science, where natural signs are revitalized as valid sources of knowledge. This research emphasizes that dew ecology is a natural information system long overlooked by mainstream scientific narratives that focus excessively on numerical data while ignoring the essence of form and meaning behind the transient phenomena of dawn.

Discussion

The empirical evidence from this study suggests that dew morphology is far from a static state of matter; rather, it represents a dynamic manifestation of external kinetic forces, specifically the negotiation between water surface tension and wind-induced pressure. The observation that dewdrops transition from perfect spheres to pointed structures under wind velocities of 0.1 to 1.5 km/h, consistently orienting their apex toward the windward direction at the leaf tip, positions dew as a high-fidelity biological anemometer. This structural adaptation reveals a sophisticated fluid mechanics response that has been largely overlooked in traditional meteorological narratives. When compared to the established quantitative frameworks in dew research, such as the thermodynamic models proposed by [14], a significant epistemological gap becomes apparent. While traditional studies treat dew as a passive object of condensation and focus primarily on yield and chemical composition, this research elevates dew to an active "sign language" or natural semiotics. This shift from purely physical measurement to a "morphological ecology" allows for a deeper understanding of how micro-geometry reflects macro-climatic shifts, suggesting that the form of a droplet is as data-rich as its volume.

The predictive power of these morphological variations provides a robust argument for the revitalization of sensory-based "local epistemology" within modern environmental science. The presence of "sand dew" and "minute dew" on specific leaf regions serves as a reliable precursor to extreme weather transitions that often elude localized digital instrumentation. For instance, the systematic correlation between the sudden disappearance of minute dew at 24°C and subsequent heavy rainfall within a 15 km radius demonstrates that nature provides its own early warning system through the rhythm of transience. Conversely, the transformation of sand dew amidst rising temperatures serves as an absolute indicator of forthcoming extreme heat thresholds, potentially reaching 45°C. By positioning the researcher as an immersed subject rather than a detached observer, this study argues that the qualitative "aesthetics" of dew – its neatness, positioning, and adherence – are not merely ornamental but are critical

epistemological indicators. This perspective challenges the mainstream scientific preoccupation with numerical data, suggesting that a "phenomenological lens" can achieve a level of predictive nuance that complements, and in some cases surpasses, the reach of modern meteorological instruments.

Furthermore, the integration of optical clarity into the analysis of atmospheric stability reinforces the role of dew as a reflection of the surface boundary layer's transparency. The "blurred dew" phenomenon, characterized by irregular, flattened shapes and low quantitative density, provides a direct visual readout of cloud persistence and high humidity interference. This lacks the sharp "light mirage" effect found in stable, globular dew, which typically predicts clear sky conditions and persistent heat. Unlike traditional hygrometers that provide isolated humidity percentages, the optical quality of a dewdrop functions as an integrated micro-lens that refracts the totality of current atmospheric conditions. The novelty of this research thus lies in its ability to synthesize physical morphological data with local meteorological forecasting, creating a new dimension where terms like "angel's dew" or "true dew" gain academic legitimacy. Ultimately, this study advocates for an environmental science that is more attuned to the essence of form and meaning, proving that dew ecology is a sophisticated natural information system that offers vital knowledge for environmental adaptation in an increasingly volatile climate [15].

CONCLUSION

Fundamental Finding: This research establishes that dew morphology functions as a sophisticated, dynamic natural information system that reflects broader atmospheric conditions. The study identifies a consistent correlation between droplet geometry – ranging from pointed structures under high-wind kinetic pressure to perfect globular forms in stable environments – and microclimatic shifts. These variations prove that the presence and clarity of "sand dew" and "minute dew" are not merely aesthetic occurrences but reliable epistemological indicators and precursors for short-term weather events, such as extreme heat or heavy rainfall. **Implication:** The findings advocate for a shift from traditional quantitative hydrology toward a "morphological ecology" framework, repositioning dew as a semiotic subject rather than a passive object of condensation. Practically, this study demonstrates that "local epistemology" and qualitative sensory observation remain vital sources of knowledge for environmental adaptation. By integrating aesthetic and morphological analysis into mainstream environmental science, we can achieve a more nuanced understanding of Human-Nature Interaction that complements modern meteorological instrumentation. **Limitation:** While this study successfully bridges the gap between environmental humanities and predictive science through a phenomenological lens, the current observations are primarily rooted in localized phenomenological analysis. The specific focus on "sand dew" and "minute dew" as indicators may currently be confined to the specific climatic and botanical conditions observed during the study period. **Future Research:** To further validate the universal applicability of dew morphology as a bio-indicator of global

atmospheric stability, future studies should expand these observations across diverse geographic ecosystems. Subsequent research should aim to standardize these qualitative signs into broader environmental models, ensuring that the integration of morphological analysis continues to foster a sense of completeness in our understanding of the Earth's atmospheric transitions.

ACKNOWLEDGEMENTS

The authors express their deepest gratitude to the University of Brawijaya and University Ibnu Sina for providing the institutional support and academic environment necessary to conduct this interdisciplinary research. Special appreciation is extended to the mentors and co-authors from various departments whose scientific insights were instrumental in bridging the gap between phenomenology and environmental science. We also acknowledge the local communities in the Riau Archipelago and South Sulawesi for their hospitality and assistance during the rigorous field observations at dawn. Finally, this work is dedicated to all parties who supported the documentation of these transient natural phenomena as a vital source of ecological knowledge.

REFERENCES

- [1] R. Armstrong, H. Furjan, and C. Van Auken, "Living Atmospheres," 2023.
- [2] T. Dimitriou, N. Skandalos, and D. Karamanis, "Progress in improving photovoltaics longevity," *Appl. Sci.*, vol. 14, no. 22, p. 10373, 2024.
- [3] G. Gangisetty and R. Zevenhoven, "A review of nanoparticle material coatings in passive radiative cooling systems including skylights," *Energies*, vol. 16, no. 4, p. 1975, 2023.
- [4] A. Sibley, M. Schulze, J. Jones, A. Kennedy, and C. Still, "Canopy wetting patterns and the determinants of dry season dewfall in an old growth Douglas-fir canopy," *Agric. For. Meteorol.*, vol. 323, p. 109069, 2022.
- [5] H. Li, C. Han, Y. Yang, and R. Chen, "Advances and Challenges in Dew Research on Land Surface: A Review," *Hydrology*, vol. 12, no. 12, p. 320, 2025.
- [6] D. Beysens, "Dew," *Encycl. l'Environnement [en ligne]*, 2022.
- [7] R. Ghazala, M. A. Askari, S. Guo, Z. Chen, B. Xu, and X. Wang, "Nature-inspired interface fluid behaviors, superhydrophobic material design, and droplet dynamics: A systematic review," *Nanoscale*, 2025.
- [8] D. Beysens, *The physics of dew, breath figures and dropwise condensation*, vol. 994. Springer, 2022.
- [9] A.-K. Lenz, U. Bauer, and G. D. Ruxton, "An ecological perspective on water shedding from leaves," *J. Exp. Bot.*, vol. 73, no. 4, pp. 1176–1189, 2022.
- [10] D. Beysens, S. Lefavrais, M. Muselli, P. Rébillout, and L. Royon, "A standard for dew measurement," *J. Hydrol.*, vol. 661, p. 133527, 2025.
- [11] M. Rahmati *et al.*, "Soil moisture memory: State-of-the-art and the way forward," *Rev. Geophys.*, vol. 62, no. 2, p. e2023RG000828, 2024.
- [12] M. S. Hossain, M. K. Alam, and M. S. Ali, "Phenomenological approach in the qualitative study: Data collection and saturation," *ICRRD Qual. Index Res. J.*, vol. 5, no. 2, pp. 148–172, 2024.
- [13] A. A. Alhazmi and A. Kaufmann, "Phenomenological qualitative methods applied to the analysis of cross-cultural experience in novel educational social contexts,"

- Front. Psychol.*, vol. 13, p. 785134, 2022.
- [14] F. Lobos-Roco *et al.*, "Understanding inland fog and dew dynamics for assessing potential non-rainfall water use in the Atacama," *J. Arid Environ.*, vol. 221, p. 105125, 2024.
- [15] M. Sun, W. Jiao, C. Gerlein-Safdi, M. Berkelhammer, and L. Wang, "Dew formation is associated with higher ecosystem productivity across diverse ecosystems," *Nat. Commun.*, vol. 17, p. 4211, 2026, doi: 10.1038/s41467-026-71933-8.

***Sofyang Daeng Kelana (Corresponding Author)**

Doctor Of Environmental Science, University of Brawijaya, Malang, Indonesia
Jl. MT Haryono 169, Malang 65145, Jawa Timur, Indonesia.
Email: sofyangkelana@gmail.com

Gatot Ciptadi

Faculty of Animal Husbandry, University of Brawijaya, Malang, Indonesia
Jl. Veteran, Ketawanggede, Kec. Lowokwaru, Kota Malang, Jawa Timur
Email: ciptadi@ub.ac.id

Aulanni'am

Department of Chemistry, Faculty of Mathematics and Natural Sciences,
University of Brawijaya, Malang, Indonesia,
Jl. Veteran, Ketawanggede, Kec. Lowokwaru, Kota Malang, Jawa Timur, Indonesia
Email: aulani@ub.ac.id

Bagyo Yanuwidi

Department of Biology, Faculty of Mathematics and Natural Sciences,
University of Brawijaya, Malang, Indonesia
Jl. Veteran, Ketawanggede, Kec. Lowokwaru, Kota Malang, Jawa Timur, Indonesia
Email: yanuwidi@ub.ac.id

Pattasang

University Ibnu Sina, Indonesia,
Jl. Teuku Umar, Lubuk Baja, Kota Batam, Kepulauan Riau, Indonesia
Email: pattasang.b1@gmail.com
